


In diarrhea of acute gastroenteritis...



LOMOTIL[®] Tablets Liquid

Each tablet and each 5 cc. of liquid contains:
 diphenoxylate hydrochloride 2.5 mg.
 (Warning: May be habit forming)
 atropine sulfate 0.025 mg.

• Lowers Motility • Allays Diarrhea • Limits Disability

No matter how quickly diarrhea may subside, it seldom subsides quickly enough for the patient.

The lack of laboratory methods for promptly identifying the causative organism increases the importance of symptomatic and supportive therapy.

Lomotil is a simple, highly acceptable agent, free of the major disadvantages of the opiates, for prolonging intestinal transit time and limiting the duration of diarrhea. With Lomotil to control intestinal hypermotility and diarrhea, patients are more comfortable and frequently are able to resume normal activities sooner.

Precautions: Lomotil is a federally exempt narcotic preparation of very low addictive potential. Recommended dosages should not be exceeded, and medication should be kept out of reach of children. Should accidental overdosage occur signs may include severe respiratory depression, flushing, lethargy or coma, hypotonic reflexes, nystagmus, pinpoint pupils and tachycardia; continuous observation is recommended. Lomotil

should be used with caution in patients with impaired liver function or those taking addicting drugs or barbiturates.

Side Effects: Side effects are relatively uncommon but among those reported are gastrointestinal irritation, sedation, dizziness, cutaneous manifestations, restlessness, insomnia, numbness of the extremities, headache, blurring of vision, swelling of the gums, euphoria, depression and general malaise.

**For correct therapeutic effect
 Rx correct therapeutic dosage**

Dosage: The recommended initial *daily dosages*, given in divided doses until diarrhea is controlled, are:

Children:

3-6 mo. . . ½ tsp. t.i.d. (3 mg.)
 6-12 mo. ½ tsp. q.i.d. (4 mg.)
 1-2 yr. . . ½ tsp. 5 times daily (5 mg.)
 2-5 yr. . . 1 tsp. t.i.d. (6 mg.)
 5-8 yr. . . 1 tsp. q.i.d. (8 mg.)
 8-12 yr. . 1 tsp. 5 times daily (10 mg.)

Adults: . . 2 tsp. 5 times daily (20 mg.)
 (or 2 tablets q.i.d.)

*Based on 4 cc. per teaspoonful.

Maintenance dosage may be as low as one-fourth the initial daily dosage.

SEARLE

Research in the Service of Medicine



"Yes, Doctor, the pain is gone."

'EMPIRIN'® COMPOUND with CODEINE PHOSPHATE gr. 1/2 No. 3

Each tablet contains: Codeine Phosphate gr. 1/2 (Warning—May be habit forming), Phenacetin gr. 2 1/2, Aspirin gr. 3 1/2, Caffeine gr. 1/2.

- Despite introduction of synthetic substitutes, efficacy of 'Empirin' Compound with Codeine remains unchallenged.



BURROUGHS WELLCOME & CO. (U.S.A.) INC., TUCKAHOE, N.Y.

The Bulletin of The New York Academy of Medicine, Vol. 43, No. 12, December, 1967. Published monthly by The New York Academy of Medicine, 2 East 103 Street, New York, N. Y. 10029. Entered as second-class matter February 3, 1928, at the Post Office at New York, N. Y., under the act of August 24, 1912. Postage paid at New York, N. Y. Annual subscription United States \$10.00. Canada \$11.00. All other countries \$12.00. Single copies \$2.00.

CALM

anxiety relieved...



Before prescribing, please consult complete product information, a summary of which follows:

Contraindications: Patients with known hypersensitivity to the drug.

Warnings: Caution patients about possible combined effects with alcohol and other CNS depressants. As with all CNS-acting drugs, caution patients against hazardous occupations requiring complete mental alertness (e.g., operating machinery, driving). Though physical and psychological dependence have rarely been reported on recommended doses, use caution in administering to addiction-prone individuals or those who might increase dosage; withdrawal symptoms (including convulsions), following discontinuation of the drug and similar to those seen with barbiturates, have been reported. Use of any drug in pregnancy, lactation, or in women of childbearing age requires that its potential benefits be weighed against its possible hazards.

Precautions: In the elderly and debilitated, and in children over six, limit to smallest effective dosage (initially 10 mg or less per day) to preclude ataxia or oversedation, increasing gradually as needed and tolerated. Not recommended in children under six. Though generally not recommended, if combination therapy with other psychotropics seems indicated, carefully consider individual pharmacologic effects, particularly in use of potentiating drugs such as MAO inhibitors and phenothiazines. Observe usual precautions in presence of impaired renal or hepatic function. Paradoxical reactions (e.g., excitement, stimulation and acute rage) have been reported in psychiatric patients and hyperactive aggressive children. Employ usual precautions in treatment of anxiety states with evidence of impending depression; suicidal tendencies may be present and protective measures necessary. Variable effects on blood coagulation have been reported very rarely in patients receiving the drug and oral anticoagulants; causal relationship has not been established clinically.

Adverse Reactions: Drowsiness, ataxia and confusion may occur, especially in the elderly and debilitated. These are reversible in most instances by proper dosage adjustment, but are also occasionally observed at the lower dosage ranges. In a few instances syncope has been reported. Also encountered are isolated instances of skin eruptions, edema, minor menstrual irregularities, nausea and constipation, extrapyramidal symptoms, increased and decreased libido—all infrequent and generally controlled with dosage reduction; changes in EEG patterns (low-voltage fast activity) may appear during and after treatment; blood dyscrasias (including agranulocytosis), jaundice and hepatic dysfunction have been reported occasionally, making periodic blood counts and liver-function tests advisable during protracted therapy.

Usual Daily Dosage: Individualize for maximum beneficial effects. Oral—Adults: Mild and moderate anxiety and tension, 5 or 10 mg t.i.d. or q.i.d.; severe states, 20 or 25 mg t.i.d. or q.i.d. Geriatric patients: 5 mg b.i.d. to q.i.d. (See Precautions.)

Supplied: Librium® (chlordiazepoxide HCl) Capsules, 5 mg, 10 mg and 25 mg—bottles of 50. Libritabs™ (chlordiazepoxide) Tablets, 5 mg, 10 mg and 25 mg—bottles of 100. With respect to clinical activity, capsules and tablets are indistinguishable.

AND CLEAR

without excessive dulling



Excessive anxiety can be an oppressive burden to the wife and mother as well as to the working man, interfering with her ability to concentrate upon the never-ending details of her daily activities and responsibilities.

For any excessively anxious individual who must constantly meet the demands of domestic or business life, Librium (chlordiazepoxide HCl) can provide dependable calming action. At the same time, Librium (chlordiazepoxide HCl), on proper maintenance dosage, does not interfere unduly with mental acuity and physical coordination, both prerequisites for adequate performance at home or on the job.

for relief of anxiety

Librium®
(chlordiazepoxide HCl)

5-mg, 10-mg, 25-mg Capsules

also available:


LIBRITABS™
(chlordiazepoxide)

5-mg, 10-mg, 25-mg Tablets

ROCHE LABORATORIES
Division of Hoffmann-La Roche Inc.
Nutley, N.J. 07110



now for your non-hospitalized patients



Serum Chemistry Graph provided is 8" x 11"

blood chemistry profiles

run on
TECHNICON®
Sequential Multiple Analyzer

only \$15⁰⁰

FOR A 12-TEST BLOOD CHEMISTRY PROFILE

Heretofore available only to patients in a few large hospitals, PATHOLOGICAL LABORATORY now provides this rapid automated analytical service for the non-hospitalized patient.

A Serum Chemistry Graph of each analysis summarizes in graphic form the results of *all 12 tests together*, superimposed over a background of normal values.

Patients may be directed to either laboratory; or blood samples can be mailed in post-paid containers provided by our laboratory. The Serum Chemistry Graph and a report in the usual numerical format will be mailed the same day samples are received. Free messenger pick-up is available in select areas.

PATHOLOGICAL LABORATORY

40-03 NATIONAL ST., CORONA, N.Y. 11368

TELEPHONE 212-639-5403

Also available through JERICHO-BIRCHWOOD LABORATORY, 59 Birchwood Park Drive, Jericho, N.Y.

TELEPHONE 516-938-0900

**send for
post-paid
blood sample
mailing
containers**

PATHOLOGICAL LABORATORY • 40-03 National St., Corona, N.Y. 11368

Gentlemen:

- ☐ Please send additional literature
- ☐ Please send post-paid containers for blood samples

Dr.

Street

City State Zip



An infinitely precious gift

*"Health and good estate of body are
above all gold, and a strong body above
infinite wealth."—Ecclesiasticus 30: 15*

Good health is the gift besides which all others are insignificant.

It is our business at GHI to pay for the care that you, as a physician, provide to make and keep your patients healthy. We extend to the medical profession and to subscribers, past, present, and future, the wish that the best of health may be theirs.



Group Health Insurance, Inc. / 221 Park Avenue South / New York

Maybe you don't want
your patients
to halve Hygroton®.
chlorthalidone



Here's the Hygroton they don't have to halve

Indications: Hypertension and many types of edema involving retention of salt and water.

Contraindications: Hypersensitivity and most cases of severe renal or hepatic disease.

Warning: With the administration of enteric-coated potassium supplements, which should be used only when adequate dietary supplementation is not practical, the possibility of small bowel lesions (obstruction, hemorrhage, and perforation) should be kept in mind. Surgery for these lesions has frequently been required and deaths have occurred. Discontinue enteric-coated potassium supplements immediately if abdominal pain, distention, nausea, vomiting, or gastrointestinal bleeding occur.

Use with caution in pregnant patients, since the drug may cross the placental barrier and adverse reactions which may occur in the adult (thrombocytopenia, hyperbilirubinemia, altered carbohydrate metabolism, etc.) are potential problems in the newborn.

Precautions: Antihypertensive therapy with Hygroton should always be initiated cautiously in postsympathectomy patients and in patients receiving ganglionic blocking agents or other potent antihypertensive drugs, or curare. Reduce dosage of concomitant antihypertensive agents by at least one-half. Barbiturates, narcotics or alcohol may potentiate hypotension. Because of the possibility of progression of renal damage, periodic determination of the BUN is indicated. Discontinue if the BUN rises or liver dysfunction is aggravated. Hepatic coma may be precipitated.

Electrolyte imbalance, sodium and/or potassium depletion may occur. If potassium depletion should occur during therapy, Hygroton should be discontinued and potassium supplements given, provided the patient does not have marked oliguria.


Take special care in cirrhosis or severe ischemic heart disease and in patients receiving corticosteroids, ACTH, or digitalis. Salt restriction is not recommended.

Adverse Reactions: Nausea, gastric irritation, vomiting, anorexia, constipation and cramping, dizziness, weakness, restlessness, hyperglycemia, hyperuricemia, headache, muscle cramps, orthostatic hypotension, aplastic anemia, leukopenia, thrombocytopenia, agranulocytosis, impotence, dysuria, transient myopia, skin rashes, urticaria, purpura, necrotizing angitis, acute gout, and pancreatitis when epigastric pain or unexplained G.I. symptoms develop after prolonged administration. Other reactions reported with this class of compounds include: jaundice, xanthopsia, paresthesia, and photosensitization.

Average Dosage: One tablet with breakfast daily or every other day.

Availability: White, single-scored tablets of 100 mg. and aqua tablets of 50 mg., in bottles of 100 and 1000. (B)46-230-D

For full details, please see the complete prescribing information.



New Hygroton 50 mg. from Geigy

to go with the Hygroton 100 mg. you know.



Geigy Pharmaceuticals
Division of
Geigy Chemical Corporation
Ardsley, New York

easy does it!

tear, moisten, compare—that's all!



Over 4½ million prescriptions already filled at Macy's

*All 12 Macy Prescription Centres
satisfy your patients with
efficient professional service*



- You may phone in your patients' prescriptions (except for narcotics, amphetamines, and barbiturates).
- Your patients may charge their prescriptions on their Macy Shopping Accounts or C-T Plans
- Your patients may have their prescriptions delivered anywhere free of extra charge.
- We serve your patient with a full line of convalescent and diabetic needs.

Macy's Herald Square, OX 5-4400
Macy's Parkchester, TA 8-7000
Macy's Jamaica, OL 7-9000
Macy's Flatbush, UL 6-5000
Macy's Roosevelt Field, PI 6-8200
Macy's Huntington, AR 1-3000
Macy's White Plains, WH 6-5015
Macy's New Haven, Conn., 624-9271
Macy's Bay Shore, MO 5-8400
Macy's Queens, AR 1-9100
Macy's Colonie, 459-1950
Macy's New Rochelle, 633-7700

BULLETIN of THE NEW YORK ACADEMY OF MEDICINE

Published Monthly

Annual subscription price
\$10.00 in the United States
Canada, \$11.00
All other countries, \$12.00
Single Copies \$2.00

The Bulletin contains scientific papers presented at the Academy, including:

Stated and Annual Meeting Addresses

Academy Section Papers

Panel Meetings and Symposia

New York Pathological Society—Abstracts

Televised Clinical Science Seminars

Public Health Reports

and Other Original Contributions

LIBRARY BIND YOUR MEDICAL JOURNALS

"Your library deserves it"

Free Pick-Up and Delivery

Colors Matched

DALEE BOOK CO.

1359 EAST 38th STREET
BROOKLYN, N. Y. 11234

ES 7 - 0250

(201) - 792 - 1222

A PRESTIGE OFFICE IN MANHATTAN

ON A CONVENIENT, ECONOMICAL RENTAL PLAN

PAY ONLY FOR THE TIME & SERVICES YOU USE!!!

at the

Lexington Professional Center

WHERE

A distinguished professional building at a prestige address . . . 133 East 73 St. (between Park & Lexington Aves.) . . . fully equipped, handsomely furnished, air-conditioned.

SERVICES

Everything you expect of a modern office . . . receptionists, telephone answering, secretaries, private office nurses . . . and more. Some services are extra, but are paid for only when used.

FACILITIES

Everything you expect of a well-equipped medical center, from pharmacy to radiology department. A large lounge and conference room is available at all times without charge.

ADVANTAGES

The ideal rental plan if you use your office just a few hours a day, a week, or all day, or every day . . . and the services you need are always available.

ECONOMY

The costs are phenomenally little since you pay only for the hours and facilities you use . . . at a low hourly rate plus a nominal monthly fee. NO CAPITAL INVESTMENT, no office to manage.

SUITES ALSO AVAILABLE FOR FULL TIME RENTAL

133 East 73 Street
New York, N. Y. 10021



212-UN 1-9000
Mrs. R. Freund, Exec. Dir.

Lexington Professional Center

STANDARD WORKS**Reprinted by HAFNER Publishing Co.****GESELL, A. L. and Others****VISION; ITS DEVELOPMENT IN INFANT AND CHILD \$11.50****XV, 329 pages, illus. (1949) (Reprint 1967)**

"Prior to this study of Gesell, Ilg and Bullis, there have been remarkably few observations on the development of the visual functions in the infant and child, and none of the comprehensive scope that are reported in this book. The material is well presented, beautifully illustrated and organized into the main headings termed "The Growing Action System," "Developmental Optics" and "Developmental Appraisal." Each subdivision is arranged chronologically according to the age of the child. Archives of Ophthalmology, Vol. 43, 1950

KOOS, E. L.**HEALTH IN REGIONVILLE****\$6.00****177 pages, (1954) (Reprint 1967)**

The first systematic examination of what people think and why they behave as they do in regard to health.

LANGE, C. G. and JAMES, WILLIAM**THE EMOTIONS****\$5.00****135 pages, (1922) (Reprint 1967)**

The theory of emotions as organic processes has become so strongly entrenched in scientific thought that it is practically assumed today to be the basis for study of emotional life. It has also led to the development of the hypothesis of reaction or response as the basis of all mental life, a theory which is rapidly supplanting the phrenologists notion of brain. The writings of James and Lange have had a profound influence upon both their contemporary and later psychologists that the name "James-Lange Theory" has been applied to the organic theory of emotions.

Order from**STECHERT-HAFNER, INC.****31 East 10th Street, New York, N. Y. 10003**

**THE FEBRUARY 1968 ISSUE OF THE BULLETIN
WILL BE DEVOTED TO THE PUBLICATION OF PAPERS**

PRESENTED AT THE

1967 Health Conference of

The New York Academy of Medicine

***"Planning for Community Health Services:
Perspectives for Action"***

Whatever happened to the family doctor?

He's moving faster these days. But he's still a good friend as well as a medical advisor to his patients.

And whether your practice is general or specialized, when you complete your work your patients look to still another friendly quarter — skilled, knowledgeable Blue Shield — to help pay their medical bills.


This has been Blue Shield's function for more than twenty-five years—helping to maintain the confidential doctor-patient relationship in the family doctor way.

BLUE SHIELD


GREATER NEW YORK'S
United Medical Service, Inc.
Two Park Ave., New York 10016



- *ventricular tachyarrhythmias*
- *Stokes-Adams attacks*
- *acute myocardial infarction with multiple extrasystoles*
- *anesthesia-induced cardiac arrest*



a pacemaker



*Figures show
that the
best
combination for
weight control
is*

**YOUR SUPERVISION
OBEDRIN®-LA
OBEDRIN MENU PLAN**

A slim figure is the glamour goal of most women. In your practice, though, you undoubtedly see many women and men who should lose weight for fundamental health reasons. Your professional guidance plus Obedrin-LA and the Obedrin Menu Plan can help keep patients on your program longer. One tablet taken daily trickle-releases medication in a balanced ratio to curb appetite and sustain mood. Write for a free supply of the Obedrin 1000 Calorie Menu Plan.

DOSAGE: OBEDRIN-LA—1 daily, usually at 10 a.m. OBEDRIN Tablets and Capsules—1 tablet or capsule at 10 a.m. and 3 p.m. If necessary to suppress late evening hunger, another tablet or capsule may be taken at 8 p.m. OBEDRIN tablets are grooved so a half tablet can be taken if it is found sufficient for appetite control.

SUPPLY: OBEDRIN-LA—Tablets, two-layer in bottles of 50 and 250. OBEDRIN—Tablets in bottles of 100, 500 and 1000; Capsules in bottles of 100 and 1000.

Caution: Federal law prohibits dispensing without prescription.

CAUTION: Should not be given concurrently with monoamine oxidase inhibitors. It should be used with caution in patients having a sensitivity to sympathomimetic compounds or barbiturates and in cases of coronary or cardiovascular disease or severe hypertension. Excessive use of amphetamines by unstable individuals has been reported to result in a psychological dependence. In such instances, withdrawal of the medication is necessary. All medication should be used with caution in pregnant patients, especially in the first trimester.

SIDE EFFECTS: Insomnia, excitability, nervousness may occur if dosage is excessive. These occur infrequently and are mild with the recommended dosage.

"Trickle-Release" Tablets

Obedrin®-LA

Each tablet contains: Methamphetamine HCl, 12.5 mg.; Pentobarbital, 50 mg. (Barbituric Acid derivative; Warning: May be habit forming); Ascorbic Acid, 200 mg.; Thiamine Mononitrate, 1 mg.; Riboflavin, 2 mg.; Niacin, 10 mg.

Obedrin®
Tablets and Capsules

Each tablet or capsule contains: Methamphetamine HCl, 5 mg.; Pentobarbital, 20 mg. (Barbituric Acid derivative; Warning: May be habit forming); Ascorbic Acid, 100 mg.; Thiamine Mononitrate, 0.5 mg.; Riboflavin, 1 mg.; Niacin, 5 mg.

MASSENGILL

The S.E. Massengill Company • Bristol, Tenn.
New York • Chicago • Dallas • San Francisco



Tears without grief

Crying Spells...psychic tension with depressive symptoms?

"I don't know what's the matter with me lately...I cry and I cry...and I really don't know why I do."

A woman often is not conscious of the real reasons for her crying spells or refuses to admit them to herself. On probing you may find that frequent weeping, like insomnia or neurotic fatigue, often is an expression of psychic tension.

She needs sympathy and reassurance, and perhaps a calming agent to help her over her crisis. Consider prescribing Valium (diazepam) for her. It usually re-establishes calmness promptly.

Crying spells and other secondary depressive symptoms normally subside as the tension is relieved. Your patient then can cope more easily with the stresses to which she is subjected.

Valium (diazepam) is generally well tolerated, and on proper maintenance dosage usually does not impair mental acuity or ability to function. If side effects such as ataxia and drowsiness occur, they usually disappear with dosage adjustment.

Before prescribing, please consult complete product information, a summary of which follows:

Contraindications: Infants, patients with history of convulsive disorders, glaucoma or known hypersensitivity to drug.

Warning: Not of value in the treatment of psychotic patients, and should not be employed in lieu of appropriate treatment.

Precautions: Limit dosage to smallest effective amount in elderly or debilitated patients (not



more than 1 mg, one or two times daily initially) to preclude ataxia or oversedation, increasing gradually as needed or tolerated. As is true of all CNS-acting drugs, until correct maintenance dosage is established, advise patients against possibly hazardous procedures requiring complete mental alertness or physical coordination. Driving during therapy not recommended. In general, concurrent use with other psychotropic agents is not recommended. If such combination therapy is used, carefully consider individual pharmacologic effects—particularly with known compounds which may potentiate action of Valium (diazepam), such as phenothiazines, barbiturates, MAO inhibitors and other antidepressants. Advise patients against simultaneous ingestion of alcohol or other CNS depressants. Safe use in pregnancy not established. Employ usual precautions in treatment of anxiety states with evidence of impending depression; suicidal tendencies may be present and protective measures necessary. Observe usual precautions in impaired renal or hepatic function. Periodic blood counts and liver function tests advisable in long-term use.

Cease therapy gradually.

Side Effects:

Side effects (usually dose-related) are fatigue, drowsiness and ataxia. Also reported: mild nausea, dizziness, blurred vision, diplopia, headache, incontinence, slurred speech, tremor and skin rash; paradoxical reactions (excitement, depression, stimulation, sleep disturbances, acute hyperexcited states, hallucinations); changes in EEG patterns during and after drug treatment. Abrupt cessation after prolonged overdosage may produce withdrawal symptoms (convulsions, tremor, abdominal and muscle

cramps, vomiting, sweating) similar to those seen with barbiturates, meprobamate and chlordiazepoxide HCl.

Dosage—Adults: Mild to moderate psychoneurotic reactions, 2 to 5 mg b.i.d. or t.i.d.; severe psychoneurotic reactions, 5 to 10 mg t.i.d. or q.i.d.; alcoholism, 10 mg t.i.d. or q.i.d. in first 24 hours, then 5 mg t.i.d. or q.i.d. as needed; muscle spasm with cerebral palsy or athetosis, 2 to 10 mg t.i.d. or q.i.d. **Geriatric patients:** 1 or 2 mg/day initially, increase gradually as needed and tolerated. (See Precautions)

Supplied: Valium® (diazepam) Tablets, 2 mg, 5 mg and 10 mg; bottles of 50 and 500.

Roche Laboratories, Division of Hoffmann-La Roche Inc., Nutley, N.J. 07110

Valium®

(diazepam) Roche®



*useful for the relief of psychic tension
with associated depressive symptoms*

